



Comparison of motor relearning occupation-based and neurodevelopmental treatment approaches in treating patients with traumatic brain injury

***Jackson Nowa, BSc Hons (University of Zimbabwe). MSc OT (Wits).**

<http://orcid.org/0000-0002-1609-1520>

Occupational Therapist, Windhoek Central Hospital, Namibia. Postgraduate student, Department of Occupational Therapy, Dept of Occupational Therapy, School of Therapeutic Sciences, Faculty of Health Sciences, University of the Witwatersrand

Denise Franzsen, BSc OT (Wits); MSc OT (Wits; DHT (UP)); PhD (Wits).

<http://orcid.org/0000-0001-8295-6329>

Sessional Senior Lecturer, Dept. of Occupational Therapy, School of Therapeutic Sciences, Faculty of Health Sciences, University of the Witwatersrand.

Dineo Thupae, BSc OT (UWC), MPH (Wits). <https://orcid.org/0000-0001-6470-1333>

Lecturer, Department of Occupational Therapy, Dept of Occupational Therapy, School of Therapeutic Sciences, Faculty of Health Sciences, University of the Witwatersrand

ABSTRACT

Introduction: Traumatic Brain Injury (TBI) is caused by trauma related to motor vehicle accidents (MVAs), accidental falls and violence. Around the world, approximately 69 million people annually suffer a TBI due to various causes with the majority of cases affecting low and middle income countries (LMICs). The management of TBI requires a multidisciplinary approach which includes rehabilitation. The aim of the study was therefore to evaluate the outcomes of the Neurodevelopmental Treatment (NDT) and motor relearning occupation-based approaches on physical performance and self-care among adults with TBI.

Methods: An experimental research design comparing two groups was used in the study. The Fugl Meyer Assessment (FMA) was used to evaluate the motor performance and a Modified Barthel Index (MBI) was used to ascertain the functional independence of the study participants before and after interventions. Data were analysed using descriptive statistics and non-parametric tests.

Results: The motor relearning occupation-based approach showed greater improvement in the FMA total, upper extremity and wrist, and most of the MBI scores. The NDT approach showed greater improvement in pain scores.

Conclusion: The findings of the current study indicate that both motor relearning occupation-based and NDT treatment approaches show clinically significant improvement in physical performance and self-care. The study also shows that the motor relearning occupation-based approach is more preferable to the NDT for improvement of physical performance and self-care.

Key words: Self-care, Neurodevelopmental Treatment approach, Motor relearning occupation-based approach, Traumatic brain injury

INTRODUCTION

“Traumatic Brain Injury (TBI) is defined as an alteration in brain function, or other evidence of brain pathology caused by external forces”¹. Around the world, approximately 69 million people annually suffer a TBI due to various causes. The incidence of TBI has been related to low socio-economic status, and the patterns and distribution of head injury may be specific to different geographic regions/countries with African countries having higher incidents². This has been associated to how well preventive and safety measures related to occupational and road safety are implemented and enforced in each country³. Developing, low and middle-income countries (LMICs) report the highest prevalence of TBI affecting males at the age of 45 and below. Motor vehicle accidents (MVAs) account for the majority of these injuries due to poor enforcement of road and vehicle regulations^{2,4}. In countries such as South Africa, Namibia, Taiwan and India, head injury resulting from MVAs is

common for drivers, passengers and pedestrians^{2,5}. The incidence of TBI resulting from MVAs in these LMICs is reported at 56%, in comparison to the lower rate of 25% which occurs in the United States of America (USA)². Although there are no prevalence figures for TBI in Namibia where this study was conducted, reported incidences of MVAs with fatalities increased by 2% in 2016 affecting the sustainable development goals which aim at reducing road deaths and injuries by 2020⁶.

The management of TBI and acquired brain injuries requires a multidisciplinary approach which includes rehabilitation. Rehabilitation is crucial in the mitigation of the effects of impairments, activity limitations, and participation restrictions⁷ during the execution of activities of daily living (ADLs), and may result in increased dependence on others and decreased quality of life (QoL). This dependence is due to deficits in higher order structures that are involved in planning and execution of smooth coordinated movements that affect the way






Figure 1: Models and theories of motor control and their relationship to treatment approach to central nervous system (CNS) dysfunction²².

individuals execute tasks^{8,9}. The outcomes of occupational therapy for clients with TBI include a return to performance in occupations that enable them to find meaning, increase participation and satisfy their potential in life, both in their homes and their communities¹⁰.

Occupational Therapists working in neuro-rehabilitation use approaches and techniques based on neuroplasticity. These approaches reduce impairments and facilitate participation in activities in patients with TBI. However, the evidence that supports the efficacy of these techniques is limited and not conclusive especially in the management of TBI and the value of different treatment approaches has not been established. Occupational therapy practitioners more commonly use a bottom-up therapy by remediating specific sensory and motor deficits, based on neuro-facilitatory techniques such as the Neurodevelopmental Treatment approach (NDT)^{11,12}. An occupation-based, or top down approach, using a motor relearning framework that considers a holistic incorporation of the patient's everyday meaningful activities has also been proposed as supporting the return to occupational performance^{12,13}.

Therefore, the use of motor relearning occupation-based interventions and NDT approaches in the management of acute TBI was investigated to enable the development and synthesis of a body of knowledge in occupational therapy in order to determine the efficacy of the treatment approaches used in treating patients with TBI in a Namibian context.

LITERATURE REVIEW

A number of motor deficits manifest after an insult to the brain¹⁴ which subsequently affect the way individuals execute daily tasks⁸. Difficulties in mobility that are caused by problems in balance, power, coordination, and cognition are common in traumatic brain injuries¹⁵. Another devastating impairment is the loss of upper and lower extremity function (including the hand) due to paralysis/paresis¹⁶.

Motor neuro-rehabilitation is based on assumptions about the cause and nature of deficits in movement. Models which address theories of motor control related to motor learning and factors affecting motor relearning, are considered to provide the rehabilitation of motor deficits after TBI. These models and theories support the various clinical approaches, principles and techniques used by taking the concept of neuroplasticity into account¹⁷. The initial model of motor control based on the reflex theory of motor control was proposed by Sir Charles Sherrington in the 1800s. It

assumed that individuals require reflexes to perform movement and these reflexes are combined into actions that compose human behaviour¹⁸. This theory of motor control does not consider centrally generated goals, or "open-loop" control, anticipatory, nor "feed-forward," movements¹⁹. The hierarchical theory was consolidated based on the work of researchers in the early 1900s. It suggests that the central nervous system (CNS) is organised hierarchically with higher levels dominating and controlling the lower levels and cortical control of movement in a top-down approach throughout the central and peripheral nervous system^{17,18}. The implications of the use of this theory clinically when treating patients with CNS damage, is that the therapist should use a developmental sequence of movements, identify and prevent primitive reflexes while normalising tone, and facilitate 'normal' movement patterns²⁰.

The systems theory of motor control explains that the neural control over movement requires a clear understanding of body systems that are related to motion and their characteristics. This includes components of the motor programming theory of motor control and, the ecological theory developed by James Gibson in 1976 which elaborated on the interaction of the individual, the task, and the environment with the aim of eliciting a motor behaviour²⁰. Systems theory considers the human body as a mechanical system that is subject to both internal stressors such as changes in physiological states and external stressors such as gravity. A number of movements could result from interactions between external forces and a number of commands from the system can elicit different varieties of these movements. The theory tries to elaborate on how initial conditions of a system can affect the characteristics of movement¹⁸ and it incorporates neurophysiology, biomechanics, and motor learning principles. Scott Kelso & Tuller²¹ indicated that the execution of normal smooth movement is developed naturally through the practice of observable, functional occupation from a myriad of conditions and experiences. This can support techniques used in rehabilitation where environmental contexts can be modified^{17,20}. None of the theories has proven to be better than the other in explaining the regulation of motor control and movement²².

Based on theories of motor control, different approaches and techniques are used in the rehabilitation of patients with TBI (Figure 1). Most treatment approaches or neurorehabilitation protocols for motor recovery and learning are based on neuroplasticity. These neuroplasticity principles are observed after a brain injury as the

CNS connections regenerate. This results from the development of new pathways through remapping²³ and permanent changes in motor performance after continuous practice²⁴. Treatment approaches include task-oriented approaches, neuro-facilitatory techniques, virtual reality, electrical stimulation, with the most commonly used being the NDT and motor relearning approaches¹³.

Treatment approaches

The NDT approach, also known as the Bobath approach, is based on the development of reflex inhibiting postures and later reflex inhibiting patterns. NDT evolved to become a problem solving NDT approach which supported the ability of a person to maintain plasticity and to learn through challenges²⁴. The NDT approach spans all three theoretical approaches of motor control and supports the nervous system working in parallel with levels and subsystems. The basic principles of NDT include inhibitory control of abnormal movements at the same time facilitating automatic postural reactions using the therapist's hands and different techniques in goal-directed activities²⁴. The effects of NDT on TBI patients have not yet been fully explored. Research shows the efficacy on stroke patients which might be similar for patients with TBI since both conditions are acquired neurological disorders. A study conducted by Hafsteinsdottir et al²⁵ concluded that NDT was ineffective in the motor rehabilitation of stroke patients in the hospital setting. A systematic literature search conducted by Kollen et al²⁶ showed that overall, this approach is not superior to alternative approaches. Díaz-Arribas et al²⁷ showed moderate proof for greater results of alternative approaches in motor control and dexterity in the upper extremity.

The motor relearning approach (which includes the task-oriented approach and occupation-based approach) was a product of work by Carr and Shepherd²⁸ which assumes that the brain is dynamic and capable of organising itself after injury or insult. The approach is task-oriented because it encourages the use of meaningful activities that are contextually based and incorporates active participation to achieve functional recovery and motor relearning by repetitive and intensive practice²⁹. Although research on the effect of this approach is limited in patients with TBI, studies show the motor relearning programme has significant effects on functional outcomes and rehabilitation of patients with stroke.

A study conducted by Chan et al³⁰ used a matched-pair randomised control trial with 52 outpatients who suffered a cerebrovascular accident and found the motor relearning programme to be more effective in enabling functional recovery of these patients. However, they stated that both conventional and function-based activities should be implemented in neuro-rehabilitation. Similar findings were reported by Immadi et al³¹ whose study revealed the efficacy of a repetitive task practice motor relearning programme compared to other conventional physiotherapy treatments.

Research which compared the effect of a motor relearning and NDT approach on patients with stroke by Langhammer and Stanghelle³² indicated that patients who received motor relearning therapy had early hospital discharge, with greater improvement in motor function and ADLs than those treated with NDT. Chan et al³⁰ agree that patients treated three months' post-stroke with the motor relearning strategies have more favourable outcomes in self-care and execution of ADLs and they showed a better transfer of skills learned to other occupations. However, the intervention did not have an effect on balance, speed or outdoor mobility³⁰. A study

by Krutulyte et al³³ on 240 participants who have suffered a stroke, showed that task-oriented therapy in a motor relearning programme was preferred, but there is not enough evidence supporting the use of this approach over the others²⁶. A Cochrane review which covered four studies on the motor relearning approach indicated that interventions did not show a higher significant clinical effectiveness from other conventional neurorehabilitation approaches³⁴.

Evidence of the efficacy of the approaches used in occupational therapy to improve treatment outcomes and provide treatment programmes that are cost-effective and have positive effects on occupational performance is therefore required.

AIM OF THE STUDY

This study determined the outcomes of the NDT and motor relearning occupation-based approach on physical performance and self-care among adults post-acute TBI and compared the results of the two approaches in a Namibian setting.

METHODS

An experimental research design, comparing two groups with a pretest-posttest assessment was used in the study³⁵. This is a design with two treatment groups where participants were assigned randomly to the groups, to consider the difference in treatment approaches on each group's participants³⁶. No control group was included as all participants with TBI were receiving treatment using either a motor relearning occupation-based or NDT approach. Pretest-posttest study designs are mostly used with experimental research designs because they are useful in assessing change in variables over time which can be used to compare two or more groups. The difference between interventions can be used to compare the effectiveness of treatment approaches.

The participants were recruited from the Katutura Intermediate Hospital in Windhoek, a tertiary institution which serves as a referral hospital for all the regions in Namibia. The hospital has an 880-bed capacity, and the occupational therapy department caters to most of the wards including the neurology and internal medicine wards to which patients with TBI are admitted. The patients from these wards usually spend 12 weeks on average in the specialised wards to allow for their stabilisation, early intervention, and rehabilitation before discharge.

Approximately 63 patients with TBI were admitted to the hospital per month between the period November 2017 and April 2018. Based on a difference of 11 points with an SD of 15 on the MBI between the groups, set at a significance of 0.05 and over six months, the confidence interval of 15 and a power of 80%, a sample of 30 participants per group were used in the study³². Inclusion criteria used were adults aged eighteen years and above with mild to moderate TBI (GCS Score 9 - 15) with evidence of decreased level of consciousness on admission and a present Glasgow Coma Scale (GCS) of 15/15. Patients were recruited if they had motor or sensory dysfunction in at least one limb.

Instrumentation and Outcome measures

Demographic questionnaire

The researcher developed a demographic questionnaire to determine the demographic characteristics of the study participants which assisted in the description of participants and to better understand their context for better analysis of the data. The demographic details included sex, age, level of education, marital status, occupation, cause of injury and the participants' GCS.

Fugl Meyer Assessment of Physical Performance (FMA)

The FMA is an instrument used to evaluate motor performance after a neurological lesion. It evaluates six categories of the patient; the amount of movement, pain, sensitivity, motor function of the upper limb and lower limb, balance, coordination, and speed^{37,38}. The scale adds up to 100 points for normal movement; 66 for upper limb and 34 for lower limb^{39,40}.

A score of 0 shows that there is no movement observed, 1 shows that the movement is minimal and 2 shows that a full range of movement has been achieved³⁷. A change in the score of 4.25 to 7.25 is seen as a clinically important difference (CID). A Global Rating Scale of Change (GROC) for the FMA indicates a change of > 50% is excellent, a change of 30% -50% is marked, 30% -10% is moderate and < 10% is slight⁴¹.

Modified Barthel Index (MBI)

The modified Barthel Index is a measure of functional independence in patients who have suffered a lesion in their brains. It provides objective and quantifiable measures of a patient's functioning. The MBI five-point scoring system shows the level of ability in self-care and their clinical status. Items are scored from 0-15, 0-10 and 0-5, a score of 99 shows "slight dependence", a score below 90 shows "moderate dependence", a score below 60 describes "severe dependence" and a score below 20 indicates "total dependence"⁴².

Research Procedure

Ethical clearance to conduct the study was obtained from the Human Research Ethics Committee (HREC) at the University of Witwatersrand (ethical clearance number: M180970). Permission to conduct the study at Katutura Intermediate Hospital was obtained from the Ministry of Health and Social Services in Namibia through the Office of the Medical Superintendents at the hospital. Patients were invited to participate in the study if they met the inclusion criteria for the study. Informed consent from the participants was sought. Family members and guardians of vulnerable participants with a cognitive ability at Rancho Los Amigos Scale Level VII and below signed informed consent on the participants' behalf.

An occupational therapist research assistant was responsible for the random assignment of participants into the two treatment groups, with 30 allocated randomly to a motor relearning occupation-based group and 30 allocated to the NDT group using a random numbers table.

Another occupational therapist performed a pre-test assessment using the FMA of physical performance and the MBI on all the patients recruited into the study. The intervention using the two approaches was carried out in the occupational therapy department where participants were seen by two different occupational therapists. Therapists treated patients in different areas and the researcher who was completing a postgraduate course in neurosciences was involved with Group 1 – motor relearning occupation-based approach. A second occupational therapist with a postgraduate qualification in NDT was responsible for Group 2 – NDT approach. The motor relearning occupation-based programme used in the current study involved occupation specific training in a hospital milieu according to task demands. Training of performance skills and patterns were required for the particular tasks chosen by participants. The therapists did not follow developmental sequences and progression was achieved by increasing the complexity of the task. Therapist and patients both participated in analysis and correction of the movements for completion of tasks and emphasis was placed on repetition⁴³. The therapist in the NDT

group focused on training of normal movement patterns, normal postures and isolated weight shift during movement. Emphasis was put on testing, training of response to handling, protective reactions, postural control, and equilibrium reactions without task-specific movement patterns. The guidelines employed focused on developmental sequences and movements were elicited in prone, supine, sitting, standing and walking. The therapist analysed and corrected the movements then the participants had to follow the guidelines given by the therapist. The main guidelines included influencing of spasticity, avoidance of abnormal patterns of movement and facilitation of normal movement patterns⁴⁴.

Following the routine practice of the occupational therapy department, one-hour daily interventions, five days a week for a period of four weeks were implemented with each participant until 20 sessions had been recorded. Participants who were discharged continued treatment as outpatients in their respective groups until 20 sessions were completed. They were provided with transport fares to attend occupational therapy as outpatients. After 20 treatment sessions were completed, a post-test assessment was conducted by the occupational therapist who completed the pre-test assessment using the Fugl-Meyer Assessment of physical performance and the Modified Barthel Index. To prevent bias, this occupational therapist was blinded and was unaware of which therapy they were receiving³⁵.

Data Analysis

Frequencies and measures of central tendency were analysed using descriptive statistics. Inferential statistics used were non-parametric and included the Wilcoxon and Mann Whitney U test since data were ordinal and the small sample resulted in data that were not normally distributed. These statistics were used to determine within-group differences pre and post-test as well as between-group differences pre and post-test.

RESULTS

Demographics

Forty (66.7%) participants in the study were male. The participants' ages ranged from 18 to 68 years, with most participants between the ages of 25 - 34 (48.3%). These demographics did not differ significantly between the NDT and motor relearning occupation-based groups indicating that the groups were comparable for gender and age. In terms of educational level, more participants in the motor relearning occupation-based group had primary school education only.

The marital status of the participants revealed that the majority (50%) were never married, and this factor differed significantly between the NDT and motor relearning occupation-based groups ($p=0.010$) with more motor relearning occupation-based group participants cohabiting. Motor vehicle accidents accounted for the majority of participants (65%) injury, followed by violence which contributed to TBI in 30% and falls which was the cause of TBI in 5% of the participants.

Change in physical performance

Group 1- Motor relearning occupation-based approach

Upper and lower extremity

A Wilcoxon Signed Rank Test revealed a statistically significant within group increase in FMA upper extremity values for all aspects of the assessment following participation in the motor relearning group, ($p <$

Table I: Demographic Characteristics (n= 60)

Demographic characteristics		Total group	Neuro-developmental group	Motor relearning occupation-based group	p value
		n(%)			
Sex	Male	40(66.7)	19(63.3)	21(70.0)	0.421
	Female	20(33.3)	11(36.4)	9(30.0)	
Age (Years) Mean Age (S.D) 36.5 (10.99)					0.438
	18 - 24	4(6.7)	2(6.6)	2(6.6)	
	25 - 34	29(48.3)	16(53.33)	13(43.33)	
	35 - 44	15(25.0)	6(19.9)	9(30.1)	
	45 - 54	8(13.3)	2(6.6)	6(19.9)	
	55 - 64	3(5.0)	3(9.9)		
	65+	1(1.7)	1(3.3)		
Level of Education	Primary	9(15.0)	2(6.6)	7(23.7)	0.069
	Secondary	30(50.0)	19(63.4)	11(36.6)	
	Tertiary	21(35.0)	9(30.0)	12(40.0)	
Marital Status	Cohabiting	8(13.3)		8(26.7)	0.010**
	Currently Married	21(35.0)	10(23.4)	11(36.6)	
	Divorced	1(1.7)	1(3.3)		
	Never Married	30(50.0)	19(63.3)	11(36.6)	
Current Occupation	Non-paid work (e.g. volunteer)	1(1.7)		1(3.3)	0.433
	Paid Employment	34(56.7)	19(63.4)	15(50.0)	
	Self Employed	9(15.0)	3(10)	6(19.9)	
	Student	4(6.7)	1(3.3)	3(10)	
	Unemployed	12(20.0)	7(23.4)	5(15.0)	

Significant at $p < 0.05$ * Significant at $p < 0.01$ **.

0.001) with overall total improvement was 43.9% (Table II, p45). A large effect size was found for the upper extremity, the wrist and the hand (indicating a clinical difference in movement) well as sensation and pain. Coordination values showed a medium effect size even though the median value did not change since 12 participants did show improvement. In the lower extremity, the within group increase in FMA values were all statistically significant ($p < 0.001$), with a large to medium effect sizes and a 35.2% increase in the total score.

Group 2 -Neurodevelopmental treatment approach Upper and lower extremity

A statistically significant within group increase in FMA upper extremity, wrist and hand as well as all other values following participation in NDT group, ($p < 0.001$), with a large or medium effect size. A 37.8% improvement in the total score was found. A similar statistically significant increase ($p \leq 0.001$) in all FMA lower extremity values with a total improvement of 38.3%, with large to medium effect sizes with a total improvement of 38.3% was observed for this group.

All components for both groups except coordination and passive joint motion for Group 2- NDT approach participants, achieved

the reported minimal clinically important difference (MCID) for the FMA upper and lower extremity scores.

Comparison of between-group change in physical performance

Upper extremity

The results on the Mann Whitney U test showed a significant statistical difference in total scores for Group 1 - motor relearning occupation-based approach participants and Group 2 NDT approach participants, ($p = 0.020$), with a large effect size. The difference in the scores for the upper extremity ($p = 0.014$) and wrist ($p = 0.027$) achieved significance and the medium effect sizes indicated the difference was clinically important. There was no significant difference in the change in scores for two groups for hand, coordination, sensation and passive joint motion scores. (Table IV p46).

A negative effect size and the significant difference ($p=0.010$) for upper extremity joint pain indicated that Group 2- NDT approach participants had greater improvement than Group 1 motor relearning occupation-based approach participants. Improvement for both coordination and sensation scores were greater for Group 2- NDT approach participants with small and medium effect sizes. Group 1 motor relearning approach participants had more

improvement in hand and passive joint motion scores with small effect sizes indicating little clinical significance for these results.

Lower extremity

The results for the lower extremity scores, comparing Group 1 motor relearning occupation-based approach participants and Group 2 NDT approach participants achieved no significant differences between the groups. Small effect sizes were found for all components when the groups were compared, with Group 1 motor relearning occupation-based approach participants achieving more improvement for all components.

Changes in self-care

Group 1 Motor relearning occupation-based approach

The findings revealed statistically significant increases in the Modified Barthel Index (MBI) for all ADL components ($p < 0.001$), with a large effect sizes and a total improvement on the MBI of 78.0%. (Table V, p 47)

Table II: Group 1 – within group changes using a motor relearning occupation-based approach on upper and lower extremity for physical performance

Fugl Meyer Variables	Pre-test (n=30)	Post-test (n=30)	z-value	p value	Effect size (Cohen's r)
	Median (Lower and upper quartile)	Median (Lower and upper quartile)			
Upper extremity	12.50 (10-18)	30.50 (20-36)	-4.80	≤0.001**	0.62***
Wrist	1.50 (0-5)	10.00 (7 - 10)	-4.83	≤0.001**	0.62***
Hand	1.00 (0 - 6)	10.00 (7 -12)	-4.79	≤0.001**	0.62***
Coordination	6.00 (2 - 6)	6.00 (6 - 6)	-3.23	≤0.001**	0.42**
Total	24 (10 – 33)	52.00 (46 – 62)	-4.62	≤0.001**	0.60***
Sensation	6.00 (6 - 12)	12.00 (10 - 12)	-4.19	≤0.001**	0.54***
Passive Joint Motion	16.00 (10 - 24)	20.00 (20 - 24)	-3.64	≤0.001**	0.47**
Joint Pain	8.00 (6 - 10)	18.00 (14 - 20)	-4.74	≤0.001**	0.61***
Lower Extremity	12 (8.00 – 16.00)	24.00 (20.00 – 26.00)	-4.64	≤0.001**	0.60***
Coordination	4.00 (3.00 – 6.00)	6.00 (6.00 – 6.00)	-3.70	≤0.001**	0.48**
Total	18 (12.50 – 22.00)	30 (26.00 – 32.00)	-4.79	≤0.001**	0.62***
Sensation	6.00 (6.00 – 12.00)	12.00 (12.00 – 12.00)	-3.98	≤0.001**	0.51***
Passive Joint Motion	15.00 (10.00 – 20.00)	20.00 (20.00 – 20.00)	-3.71	≤0.001**	0.48**
Joint Pain	10.00 (7.00 – 10.50)	20.00 (18.00 – 20.00)	-4.75	≤0.001**	0.61***

Significant at p <0.05*. Significant at p <.0.01** .

Large Effect Size = 0.5***.
Medium Effect Size = 0.3**
Small Effect Size = 0.1*

Table III: Group 2 - within group changes using a neurodevelopmental treatment approach on upper and lower extremity for physical performance

Fugl Meyer Variables	Pre-test (n=30)	Post-test (n=30)	z-value	p value	Effect size (Cohen's r)
	Median (Lower and upper quartile)	Median (Lower and upper quartile)			
Upper extremity	13.50 (9.75 – 19.00)	28.00 (20.00 – 36.00)	-3.58	≤0.001**	0.46**
Wrist	3.00 (0.00 – 5.00)	7.00 (5.00 – 10.00)	-4.49	≤0.001**	0.58***
Hand	2.00 (0.00 – 7.00)	10.00 (6.00 – 14.00)	-4.64	≤0.001**	0.60***
Coordination	4.00 (3.00 – 6.00)	6.00 (6.00 – 6.00)	-3.58	≤0.001**	0.46**
Total	22.5 (12.75 – 33.5)	49 (38.75 – 63.75)	-4.79	≤0.001**	0.62***
Sensation	6.00 (6.00 – 12.00)	12.00 (10.00 -12.00)	-4.06	≤0.001**	0.54***
Passive Joint Motion	20.00 (12.00 – 24.00)	24.00 (20.00 – 24.00)	-3.53	≤0.001**	0.46**
Joint Pain	8.00 (0.00 – 12.00)	20.00 (17.25 – 24.00)	-4.65	≤0.001**	0.60***
Lower extremity	12.5 (6.00 – 16.00)	24.50 (18.00 - 26.00)	-4.71	≤0.001**	0.61***
Coordination	4.00 (2.25 – 6.00)	6.00 (6.00 – 6.00)	-3.74	≤0.001**	0.48**
Total	17.5 (6.75 – 22.00)	30.5 (24.00 – 34.00)	-4.71	≤0.001**	0.61***
Sensation	10.00 (6.00 – 12.00)	12.00 (10.00 -12.00)	-3.34	≤0.001**	0.43**
Passive Joint Motion	20.00 (10.00 – 20.00)	20.00 (20.00 – 20.00)	-3.31	≤0.001**	0.43**
Joint Pain	10.00 (6.75 – 14.50)	20.00 (15.50 – 20.00)	-4.33	≤0.001**	0.56***

Significant at p <0.05*. Significant at p <.0.01**.

Large Effect Size = 0.5***.
Medium Effect Size = 0.3**
Small Effect Size = 0.1*

Group 2 - Neuro-developmental treatment approach

The findings revealed statistically significant increase in MBI for all components with a large effect sizes and a total improvement for on the MBI of 56% (Table VI, p47).

Overall Group 1 - motor relearning occupation-based approach participants achieved a score above 60 post-test which indicated moderate independence while Group 2 NDT approach participants had a score below 60 indicating severe dependence post-test.

Comparison of between-group change in self-care

There was a significant difference in the total scores for the two groups (p = 0.002) and a medium effect size of which indicated that the Group 1 motor relearning occupation-based approach participants had more improvement in self-care with a clinically important difference (Table VII p48). Group 1 motor relearning occupation-based approach participants had a significantly larger improvement in mobility and self-care components all with large and medium effect sizes.

DISCUSSION

The demographic details were similar to a study by Samanamalee et al⁴⁵ who recorded a mean age 41.67 (SD 17.47) years and the majority of the participants being males (82%) confirming that TBI is more prevalent among young adults and males in LMICs. For occupational performance outcomes however, no significant differences were found in the current study although literature has reported that pre-injury occupation, high level of education, female sex and being married make a significant contributing factor to occupational performance outcomes¹⁴. The findings from the current study suggest that most of TBI cases were caused by vehicle-related collisions which was supported by Dewan et al² and Agrawal et al⁴³ who reported that MVAs are responsible for the silent epidemic of TBI, among the productive age groups in LMICs.

The results of the study indicate that there was a significant improvement in physical performance and self-care in both Group 1 motor relearning occupation-based approach

Table IV: Between group comparison of treatment approaches on upper and lower extremity physical performance

Physical Components	Change in Motor learning occupation-based scores	Change in Neurodevelopmental Treatment scores		
			p value	Effect size (Cohen's r)
Upper Extremity	16.00 (14.00-18.00)	13.00 (8.00-16.00)	0.014*	0.45**
Wrist	5.00 (4.00 – 10.00)	5.00 (2.00 – 5.00)	0.027*	0.40**
Hand	7.00 (5.00 – 9.00)	7.00 (4.00 – 8.00)	0.290	0.19*
Coordination	0.00 (0.00 – 3.00)	1.00 (0.00 – 3.00)	0.970	-0.01
Total	29.00 (25.00 – 34.00)	25.00 (20.00 – 30.00)	0.020*	0.42**
Sensation	4.00 (0.00 – 6.00)	4.00 (0.00 – 6.00)	0.864	-0.03
Passive joint motion	5.00 (0.00 – 10.00)	2.00 (0.00 – 8.00)	0.468	0.13*
Joint pain	10.00 (6.00 – 10.00)	12.00 (8.00 – 14.00)	0.010**	-0.47**
Lower Extremity	12.00 (8.00 – 18.00)	10.00 (8.00 – 16.00)	0.482	0.13*
Coordination	2.00 (0.00 – 3.00)	1.00 (0.00 -3.00)	0.848	0.02
Total	12.50 (8.75 – 19.25)	13.00 (8.00 – 19.25)	0.711	0.05
Sensation	4.00 (0.00 – 6.00)	0 (0.00 – 4.50)	0.131	0.20*
Passive joint motion	4.00 (0.00 – 10.00)	0.00 (0.00 – 10.00)	0.862	0.02
Joint pain	10.00 (6.00 – 10.00)	9.50 (3.50 – 10.00)	0.197	0.17*

Significant at $p < 0.05^*$. Significant at $p < 0.01^{**}$.

Large Effect Size = 0.5^{***} .
 Medium Effect Size = 0.3^{**}
 Small Effect Size = 0.1^*

participants and Group 2 NDT approach participants. The percentage improvement in the physical performance of the upper extremity in Group 1 was greater at 43.9% compared to the lower extremity at 35.2% in the current study. The notable improvements in upper extremity as compared to lower extremity found in the current study could be attributed to the fact that occupational therapists tend to focus more on the upper extremity than the lower extremity as suggested by Rowland et al⁴⁶.

The participants in the motor relearning occupation-based group were found to have a 78% improvement on the MBI indicating the effectiveness of this approach in self-care in adults with TBI. This was confirmed by statistically significant results and the observed effect sizes which were high, describing the high clinical importance of the approach. It appears that patients using motor relearning concepts in an occupation based programme regain function and independence by being involved in occupations they find meaningful to them since these occupations improve cortical representation of their skill sets⁴⁷. Occupation-based treatments done in a hospital setup that mimic the home environment improved neuroplasticity, increased functional use of the affected upper and lower extremities, and improved occupational performance⁴⁷. In the current study observations were made that participants using the motor relearning occupation-based group made more efforts in fulfilling their occupations

as an end goal. This was also noted by Giuffrida et al⁴⁸ who stated that a significant improvement in performance is seen more in random practice than in structured practice and a transfer of skill is noted in the latter.

In this study, the motor relearning occupation-based approach was thus found to be effective in enhancing physical performance in the upper extremity in particular and task performance after TBI. Similar findings were noted in a number of other studies on patients with stroke. Chan et al³⁰ found that the patients' recovery was noted by significant improvement in physical ability in balance as well as for all aspects of self-care assessed by the Functional Independence Measure. Kollen et al²⁶ also concluded that activities when used in inpatient therapy can elicit functional recovery when the activities are relatively challenging to the individual performing the task. The studies by Kollen et al²⁶ and Chan et al³⁰ found that MBI (as in the current study) showed responsiveness for improvements in transfers, bathing, personal hygiene, dressing and feeding. Although the study by Kollen et al²⁶ provided evidence supporting the lack of superiority of the NDT approach in managing sensorimotor deficits in the upper extremity and the lower extremity as well as in execution of ADLs, in the current study there was a significant improvement in physical performance and self-care for participants in the NDT approach participants.

There are some neurological changes that

are expected to occur due to a brain lesion that affect motor pathways and connections, these include loss of power, differences in tone and poor communication with the cortical areas that affect movement⁴⁹. These symptoms can be addressed by using NDT which focuses mainly on the motor units and the physical performance domain assessed by the FMA showed significant improvements in upper and lower extremity total scores, sensation, coordination, movements and pain scores in the current study. The percentage improvement in the physical performance of the upper extremity in Group 2 was slightly lower at 37.8% compared to the lower extremity at 38.2% in the current study. The improvement seen in the upper and lower extremity in Group 2 was similar as therapy applies equally to both extremities since the approach supports clinical reasoning which allows the therapist to focus on individual deficits. In the current study, the therapist in the NDT group focused on training of normal movement patterns, normal postures and isolated weight shift during movement²⁴.

No published studies on the effectiveness of NDT for adult TBI patients were sourced but Huseysinoglu et al⁵⁰ concluded that participants treated with an NDT approach showed significant improvement in physical performance including sensorimotor function, quality and speed of movement in paretic upper extremity after stroke. They did not indicate the effect of treatment in the lower extremity. A study by Bhalariao et al⁴⁴ however revealed that post-therapy participants treated with an NDT approach showed significant improvement in both upper and lower extremities on all scales of motor function and functional mobility after stroke.

The improvement seen in self-care for Group 2 participants supports the hierarchical approach in NDT that follows steps that need to be taken to achieve

Table V: Group 1 – within group changes using a motor relearning occupation-based approach for self-care

Barthel Index	Pre-test (Md) (n=30)	Post-test (Md) (n=30)	z-value	p value	Effect size (Cohen's r)
	Median (Lower and upper quartile)	Median (Lower and upper quartile)			
Chair/bed Transfers	0.00 (0.00 – 3.00)	12.00 (8.00 – 12.00)	-4.82	≤0.001**	0.62***
Ambulation	1.50 (0.00 – 3.00)	8.00 (8.00 – 12.00)	-4.82	≤0.001**	0.62***
Ambulation/ Wheel-chair	1.00 (0.00 – 1.00)	4.00 (3.00 – 5.00)	-4.87	≤0.001**	0.62***
Stair Climbing	0.00 (0.00 – 0.00)	5.00 (2.00 – 8.00)	-4.59	≤0.001**	0.59***
Toilet Transfers	1.00 (0.00 – 2.00)	8.00 (5.00 – 10.00)	-4.83	≤0.001**	0.62***
Bowel control	5.00 (3.75 – 10.00)	10.00 (5.00 – 10.00)	-3.56	≤0.001**	0.46**
Bladder control	5.00 (3.75 – 10.00)	10.00 (5.00 – 10.00)	-3.56	≤0.001**	0.46**
Bathing	0.00 (0.00 – 1.50)	4.00 (3.00 – 5.00)	-4.81	≤0.001**	0.62***
Dressing	0.00 (0.00 – 2.00)	8.00 (5.00 – 8.00)	-4.83	≤0.001**	0.62***
Personal hygiene	0.00 (0.00 – 1.00)	4.00 (3.00 -5.00)	-4.83	≤0.001**	0.62***
Feeding	2.00 (0.00 – 2.00)	8.00 (5.00 – 10.00)	-4.83	≤0.001**	0.62***
Total	2.00 (10.50 – 34.00)	80.00 (59.50 – 86.50)	-4.78	≤0.001**	0.62***

Significant at p <0.05*. Significant at p <.0.01**.

Large Effect Size = 0.5***.
Medium Effect Size = 0.3**
Small Effect Size = 0.1*

Table VI: Group 2 – within group changes using a neurodevelopmental treatment approach for self-care

Barthel Index	Pre-test (Md) (n=30)	Post-test (Md) (n=30)	z-value	p value	Effect size (Cohen's r)
	Median (Lower and upper quartile)	Median (Lower and upper quartile)			
Chair/bed Transfers	0.00 (0.00 – 3.00)	8.00 (3.00 -12.00)	-4.83	≤0.001**	0.62***
Ambulation	0.00 (0.00 – 0.00)	5.50 (3.00 – 12.00)	-4.78	≤0.001**	0.62***
Ambulation/ Wheel-chair	0.00 (0.00 – 0.00)	3.50 (3.00 – 4.00)	-4.86	≤0.001**	0.63***
Stair Climbing	0.00 (0.00 – 0.00)	2.00 (0.00 – 2.75)	-4.16	≤0.001**	0.54***
Toilet Transfers	0.00 (0.00 – 0.00)	5.00 (2.00 – 8.00)	-4.83	≤0.001**	0.62***
Bowel control	0.00 (0.00 – 5.00)	10.00 (5.00 – 10.00)	-4.33	≤0.001**	0.56***
Bladder control	0.00 (0.00 – 5.00)	10.00 (5.00 – 10.00)	-4.26	≤0.001**	0.55***
Bathing	0.00 (0.00 – 0.00)	3.00 (3.00 – 3.00)	-4.94	≤0.001**	0.64***
Dressing	0.00 (0.00 – 0.00)	5.00 (2.00 – 5.00)	-4.87	≤0.001**	0.63***
Personal hygiene	0.00 (0.00 – 0.00)	3.00 (3.00 – 4.00)	-4.82	≤0.001**	0.62***
Feeding	0.00 (0.00 – 2.00)	5.00 (2.00 – 10.00)	-4.81	≤0.001**	0.62***
Total	2.00 (0.00 -12.75)	58.00 (38.50 – 72.25)	-4.78	≤0.001**	0.62***

Significant at p <0.05*. Significant at p <.0.01**.

Large Effect Size = 0.5***.
Medium Effect Size = 0.3**
Small Effect Size = 0.1*

functional recovery by eliciting normal movements and preventing compensation²⁴. The NDT approach is a bottom-up approach which relies on treating underlying symptoms with the assumption that this will lead to an improvement in occupational performance. The findings of the current study were very different from Hafsteinsdóttir et al²⁵ which found NDT ineffective as a treatment modality for self-care in stroke patients. It can be assumed in the current study that more emphasis was placed on self-management for participants in Group 2, since the therapist involved in the NDT programme was an occupational therapist. She may well have placed more emphasis on participation in self-care since there was a change of over 50% in the self-care assessed on the MBI which was higher than the change seen in the physical performance for the upper and lower extremity.

A study by Lannin and McCluskey¹¹ stated that there was no comprehensive evidence of effects of different treatment approaches used in TBI, however this was not to be mistaken for no evidence of efficacy. In the current study, between-group comparison showed that most components for both groups achieved the reported minimal clinically important difference (MCID) for the FMA upper extremity scores. Although the MCID included in the current study were recorded for stroke patients, it was assumed this could be applied to patients with TBI since they are all acquired brain injuries⁵¹.

Overall, the change for the upper extremity was higher in Group 1 - motor relearning occupation-based approach participants was significantly higher with a total percentage increase of 43.9% compared to 37.8% in Group 2 - NDT approach participants. The

Table VII: Comparison of treatment approaches on self-care

Barthel index	Change in Motor learning occupation-based scores(n = 30)	Change in Neurodevelopmental Treatment scores (n = 30)	p value	Effect size – Cohen’s r value
	Median (lower and upper quartile)	Median (lower and upper quartile)		
Chair/bed transfers	12.00 (8.00 – 12.00)	8.00 (3.00 – 12.00)	0.002**	0.39**
Ambulation	8.00 (8.00 – 12.00)	5.50 (3.00 – 12.00)	0.072	0.23*
Ambulation/wheelchair	4.00 (3.00 – 5.00)	3.50 (3.00 – 4.00)	0.035*	0.28*
Stair climbing	5.00 (2.00 – 8.00)	2.00 (0.00 -2.75)	0.027*	0.31**
Toilet transfers	8.00 (5.00 – 10.00)	5.00 (2.00 – 8.00)	0.023*	0.31**
Bowel control	10.00 (5.00 – 10.00)	10.00 (5.00 – 10.00)	0.413	0.10*
Bladder control	10.00 (5.00 – 10.00)	10.00 (5.00 – 10.00)	0.364	0.12*
Bathing	4.00 (3.00 – 5.00)	3.00 (3.00 – 3.00)	0.001**	0.53***
Dressing	8.00 (5.00 – 8.00)	5.00 (2.00 – 5.00)	0.001**	0.43**
Personal Hygiene	4.00 (3.00 – 5.00)	3.00 (3.00 – 4.00)	0.008**	0.34**
Feeding	8.00 (5.00 – 10.00)	5.00 (2.00 – 10.00)	0.096	0.22*
Total	80.00 (59.50 – 86.50)	58.00 (38.50 – 72.25)	0.004	0.37**

Significant at $p < 0.05^*$. Significant at $p < 0.01^{**}$.

Large Effect Size = 0.5^{***} .
 Medium Effect Size = 0.3^{**}
 Small Effect Size = 0.1^*

results from the current study are in line with a study by Langhammer and Stanghelle³² who concluded that treatment that used the motor relearning approach was preferred in improving upper extremity physical performance to the one using the NDT approach in the acute rehabilitation of stroke patients. Skubik-Peplaski et al⁴⁷ also showed a significant improvement in total FMA scores in occupation-based intervention programme.

However, the current study illustrated that the change in participants in Group 1 was not consistently better than that for participants in Group 2. While there was significantly greater change for the upper extremity and wrist on the FMA for Group 1, there was no significant difference between the two groups in hand, passive joint movement, sensation and coordination. However, the effect size favoured Group 1 - motor relearning occupation-based ap-

proach for improvement in hand and passive joint movement while co-ordination was favoured in Group 2. These findings are supported by Platz et al⁵² who report the efficacy of specific techniques in hand rehabilitation is not proven for patients with TBI, who have recovery of other upper extremity function. They found continued limitations in hand function including reduced speed and accuracy affecting coordination after discharge from in-patient rehabilitation. Since optimum recovery was seen two years post injury the researcher assumed that four weeks of rehabilitation post injury were too early to find noticeable changes in the both groups in the current study in terms of hand functioning and coordination^{53,54}.

There was no difference found for sensation between the groups even though the NDT approach used some sensory input through positioning as part of the treatment.

This could be attributed to the length of time for which the participants received input in the treatment approach which was not enough to cause significant changes in the client factors such as sensation. On the other hand, the participants in Group 2 - NDT approach had significantly more improvement for pain scores compared to the participants in Group 1 - motor relearning occupation-based approach. This is likely attributed to the use of handling and positioning that NDT focuses on, Walsh⁵⁵ stated that poor positioning exacerbates shoulder pain and other types of pain in patients with stroke. This can be addressed directly when using an NDT approach.

When comparing the differences in self-care between the groups, Group 1 - motor relearning occupation-based approach participants also had significantly more improvement in MBI scores than Group 2 - NDT approach participants. Medium to large effect sizes were found for seven of the domains of the MBI indicating important clinical difference for the two groups. A significant improvement noted in self-care domains was noted for Group 1 for ambulation/wheelchair, transfers, bathing, stair climbing, personal hygiene, dressing, and feeding when using the motor relearning occupation-based approaches. This was confirmed by the much higher percentage change in self-care assessed on the MBI (78%) for participants in Group 1 - motor relearning occupation-based approach than the participants in Group 2 - NDT approach (56%). This is important since Zhu et al⁵⁶ have shown that self-care scores are a better predictor of recovery from moderate TBI at one year than age and GCS.

The results of the current study appear to support that the use of the motor relearning occupation-based approach for outcomes in physical performance in the upper limb and in self-care compared to the NDT approach. Since there was no significant difference for the lower extremity, the findings of Seneviratne and Reimer⁵⁷, who concluded, when comparing the NDT and the motor relearning approach that mixed conclusions may be found was accepted. They are in agreement with some other studies that indicate that a motor relearning occupation-based approach should be added to the current occupational therapy theory and practice and this approach can be considered complimentary to NDT rather than superior to it. Therefore, the use of both approaches for various goals in occupational therapy should be considered.

Limitations

Even though the sample size was relatively small, data gathered can be generalised to occupational therapy practice in Namibia since the results still provide meaningful findings and insights. There may have been unknown confounding variables such as the expertise that the therapists had on the treatment approaches and the motivation the clients had. It is most likely that the patients who showed improvements put in more effort than the other clients, there was no way to measure the effort put in by participants. All participants received other forms of therapy during this period and it is not clear what effect this may have had on their improvement. It is possible that some aspects of the individual therapists and the environment such as a treatment setting, ways of instruction and feedback might have led to some biases that were not controlled for in this study which might have affected the effect sizes.

CONCLUSION

The findings of the current study indicate that both motor relearning occupation-based and NDT treatment approaches are effective in occupational therapy treatment of acute traumatic brain injuries and there was a significant improvement in physical performance as well as self-care. However, the motor relearning occupation-based approach was found to be significantly superior in self-care outcomes as well as some upper extremity outcomes. No difference was found for lower extremity outcomes between the two treatment approaches. The NDT treatment approach was found to be significantly superior in addressing joint pain domains which had a significant increase compared to the other group.

REFERENCES

1. Menon DK, Schwab K, Wright DW, Maas AI. Position statement: Definition of traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*. Elsevier Inc. 2010; 91(11): 1637–1640. <https://doi.org/10.1016/j.apmr.2010.05.017>
2. Dewan MC, Rattani A, Gupta S, Baticulon RE, Hung Y-C, Maria P, et al. Estimating the global incidence of traumatic brain injury. *Journal of Neurosurgery*. Journal of Neurosurgery. 2018; 4 (2018): 1080-1097. <https://doi.org/10.3171/2017.10.JNS17352>.
3. Zolotor AJ, Runyan DK, Shanahan M, Durrance CP, Nocera M, Sullivan K, et al. Effectiveness of a statewide abusive head trauma prevention program in North Carolina. *JAMA Pediatrics*. American Medical Association. 2015; 169(12): 1126–1131. <https://doi.org/10.1001/jamapediatrics.2015.2690>
4. Bustamante VK, Montes S, Madrigal MDJ, Burciaga A, Martínez ME, Johnson MJ. Technology-assisted stroke rehabilitation in Mexico: A pilot randomized trial comparing traditional therapy to circuit training in a Robot/technology-assisted therapy gym. *Journal of NeuroEngineering and Rehabilitation*. 2016; 13(1): 1–15. <https://doi.org/10.1186/s12984-016-0190-1>
5. Naidoo D. Traumatic brain injury: The South African landscape. *South African Medical Journal*. 2013; 103(9): 613–614. <https://doi.org/10.7196/SAMJ.7325>
6. Motor Vehicle Accident Fund. Road crash and claims report 2016. 2016. Available at: https://abrie.github.io/NamOpenData2018/raw/mva_fund_road_crash_and_claims_report_2016.pdf (accessed 23.6.2019)
7. OTASA. Occupational Therapy Association of South Africa Position Statement on Rehabilitation. *South African Journal of Occupational Therapy*. 2017; 47(2): 53–55.
8. Walker WC, Marwitz JH, Kreutzer JS, Hart T, Novack TA. Occupa-

tional Categories and Return to Work After Traumatic Brain Injury: A Multicenter Study. *Archives of Physical Medicine and Rehabilitation*. 2006; 87(12): 1576–1582. <https://doi.org/10.1016/j.apmr.2006.08.335>

9. Powell JM, Rich TJ, Wise EK. Effectiveness of occupation- and activity-based interventions to improve everyday activities and social participation for people with traumatic brain injury: A systematic review. *American Journal of Occupational Therapy*. 2016; 70(3): 7003180040p1-7003180040p9. <https://doi.org/10.5014/ajot.2016.020909>
10. Soeker MS, Pape C. The Use of the Model of Occupational Self-Efficacy for Work Retraining: A Multiple Case Study. *Occupational Therapy International*. 2019; 2019(Article ID 3867816): 1-8. <https://doi.org/10.1155/2019/3867816>
11. Lannin NA, McCluskey A. A systematic review of upper limb rehabilitation for adults with traumatic brain injury. *Brain Impairment*. 2008; 9(3): 237–246. <https://doi.org/10.1375/brim.9.3.237>
12. Wolf T, Chuh J, Floyd A, McInnis T, K. Williams E. Effectiveness of Occupation-Based Interventions to Improve Areas of Occupation and Social Participation After Stroke: An Evidence-Based Review. *American Journal of Occupational Therapy*. 2015; 69(1): 11. <https://doi.org/10.5014/ajot.2015.012195>
13. Rahja M, Jolliffe L. There is moderate evidence for the effectiveness of occupation and activity-based interventions for people with traumatic brain injury although more evidence is needed for interventions provided specifically by occupational therapists. *Australian Occupational Therapy Journal*. 2018; 65(1): 73–74. <https://doi.org/10.1111/1440-1630.12449>
14. Walker WC, Pickett TC. Motor impairment after severe traumatic brain injury: A longitudinal multicenter study. *Journal of Rehabilitation Research & Development*. 2007; 44(7): 975–982. <https://doi.org/10.1682/JRRD.2006.12.0158>
15. Katz DI, White DK, Alexander MP, Klein RB, Di AK, Dk W, et al. Recovery of Ambulation After Traumatic Brain Injury. *Archives of Physical Medicine and Rehabilitation*. 2004; (6): 865-869. <https://doi.org/10.1016/j.apmr.2003.11.020>
16. Galea M, Khan F, Amatya B, Elmalik A, Klaić M, Abbott G. Implementation of a technology-assisted programme to intensify upper limb rehabilitation in neurologically impaired participants: A prospective study. *Journal of Rehabilitation Medicine*. 2016; 48(6): 522–528. <https://doi.org/10.2340/16501977-2087>
17. Cano-de-la-Cuerda R, Molero-Sánchez A, Carratalá-Tejada M, Alguacil-Diego I., Molina-Rueda F, Miangolarra-Pagea JC, et al. Theories and control models and motor learning: Clinical applications in neurorehabilitation. *Neurología (English Edition)*. SEGO; 2015; 30(1): 32–41. <https://doi.org/10.1016/j.nrleng.2011.12.012>
18. Shumway-Cook A, Woollacott HM. *Motor Control. Translating Research into Clinical Practice*. Lippincott Williams & Wilkins; Philadelphia; 2012.
19. Horak FB. Assumptions underlying motor control for neurologic rehabilitation. In *Contemporary management of motor control problems: Proceedings of the II STEP conference 1991*: 11-28. Alexandria, Foundation for Physical Therapy.
20. Roller ML, Lazaro RT, Byl NN, Umphred DA. Contemporary issues and theories of motor control, motor learning, and neuroplasticity. Chapter 3 in *Umphred's Neurological Rehabilitation - e-Book*. Lazaro RT, Reina-Guerra SG. (eds) 2013. <https://doi.org/10.1016/B978-0-323-07586-2.00013-3>
21. Scott Kelso JA, Tuller B. *A Dynamical Basis for Action Systems*. Handbook of Cognitive Neuroscienc;. 2013.

- https://doi.org/10.1007/978-1-4899-2177-2_16
22. Mathiowetz V, Haugen JB. Motor behavior research: implications for therapeutic approaches to central nervous system dysfunction. *The American Journal of Occupational Therapy*. 1994; 48(4): 733–745. <https://doi.org/10.5014/ajot.48.8.733>
 23. Chen H, Epstein J, Stern E. Neural plasticity after acquired brain injury: Evidence from functional neuroimaging. *Physical Medicine and Rehabilitation*. 2010; 2(12 SUPPL): S306–S312. <https://doi.org/10.1016/j.pmrj.2010.10.006>
 24. Raine S, Meadows L, Lynch-Ellerington M. *Bobath concept: theory and clinical practice in neurological rehabilitation*. London: John Wiley & Sons; 2013.
 25. Hafsteinsdóttir TB, Algra A, Kappelle LJ, Grypdonck MHF. Neurodevelopmental treatment after stroke: A comparative study. *Journal of Neurology, Neurosurgery and Psychiatry*. 2005; 76(6): 788–792. <https://doi.org/10.1136/jnnp.2004.042267>
 26. Kollen BJ, Lennon S, Lyons B, Wheatley-Smith L, Scheper M, Buurke JH, et al. The effectiveness of the Bobath concept in stroke rehabilitation what is the evidence? *Stroke*. 2009; 40(4): e89–e97. <https://doi.org/10.1161/STROKEAHA.108.533828>
 27. Díaz-Arribas MJ, Martín-Casas P, Cano-de-la-Cuerda R, Plaza-Manzano G. Effectiveness of the Bobath concept in the treatment of stroke: a systematic review. *Disability and Rehabilitation*. 2019; 42(12): 1636–1649. <https://doi.org/10.1080/09638288.2019.1590865>
 28. Carr HJ, Shepherd RB. *Neurological Rehabilitation - Optimizing Motor Performance*. London: Churchill Livingstone; 2011.
 29. Mulder T, Hochstenbach J. Adaptability and flexibility of the human motor system: Implications for neurological rehabilitation. *Neural Plasticity*. 2001. <https://doi.org/10.1155/NP.2001.131>
 30. Chan DYL, Chan CCH, Au DKS. Motor relearning programme for stroke patients: a randomized controlled trial. *Clinical Rehabilitation*. 2006; 20(3): 191–200. <https://doi.org/10.1191/0269215506CR9300A>
 31. Immadi KS, Achyutha KK, Reddy A, Tatakunda PK. Effectiveness of the motor relearning approach in promoting physical function of the upper limb after a stroke. *International Journal of Physiotherapy*. 2015 ;2(1): 386–390. <https://doi.org/10.15621/ijphy/2015/v2i1/60047>
 32. Langhammer B, Stanghelle JK. Bobath or motor relearning programme? A comparison of two different approaches of physiotherapy in stroke rehabilitation: A randomized controlled study. *Clinical Rehabilitation*. 2000; 14(4): 361–369. <https://doi.org/10.1191/0269215500cr338oa>
 33. Krutulyte G, Kimtys A, Krisciunas A. The effectiveness of physical therapy methods (Bobath and motor relearning program) in rehabilitation of stroke patients. *Medicina (Kaunas)*. 2003; 39(9): 889–895.
 34. Pollock A, Baer G, Campbell P, Choo PL, Forster A, Morris J, et al. Physical rehabilitation approaches for the recovery of function and mobility following stroke. *Cochrane Database of Systematic Reviews*. 2014; 45(10): 202–208. <https://doi.org/10.1002/14651858.CD001920.pub3>
 35. Kumar R. *Research Methodology. A step by step guide for beginners.. 3rd Editio*. London: SAGE Publications; 2011.
 36. Walker W. The strengths and weaknesses of research designs involving quantitative measures. *Journal of Research in Nursing*. 2005; 10(5): 571–582. <https://doi.org/10.1177/136140960501000505>
 37. Gladstone DJ, Danells CJ, Black SE. The Fugl-Meyer Assessment of Motor Recovery after Stroke: A Critical Review of Its Measurement Properties. *Neurorehabilitation and Neural Repair*. 2002; 16(3): 232–240. <https://doi.org/10.1177/154596802401105171>
 38. Fugl-Meyer A, Jaasko L, Leyman I, Olsson S, Stegling S. The Post Stroke Hemiplegic Patient. *Scandinavian Journal of Rehabilitation Medicine*. 1975; 7(2): 13–31.
 39. Padovani C, Valério C, Pires G, Pretti F, Ferreira C, Borin G, et al. Application of the Fugl-Meyer Assessment (FMA) and the Wolf Motor Function Test (WMFT) in the recovery of upper limb function in patients after chronic stroke : a literature review. *Acta Fisiatr*. 2013; 20(1): 42–49. <https://doi.org/10.5935/0104-7795.20130008>
 40. Sullivan KJ, Tilson JK, Cen SY, Rose DK, Hershberg J, Correa A, et al. Fugl-Meyer Assessment of Sensorimotor Function Standardized Training Procedure for Clinical Practice and Clinical Trials. *Stroke*. 2011; 42(2): 427–32. <https://doi.org/10.1161/STROKEAHA.110.592766>
 41. Page SJ, Fulk GD, Boyne P. Clinically Important Differences for the Upper-Extremity Fugl-Meyer Scale in People With Minimal to Moderate Impairment Due to Chronic Stroke. *Physical Therapy*. 2012; 92(6): 791–798. <https://doi.org/10.2522/ptj.20110009>
 42. Shah S, Vanclay F, Cooper B. Improving the sensitivity of the Barthel Index for stroke rehabilitation. *Journal of clinical epidemiology*. 1969; 42(8): 703–709. [https://doi.org/10.1016/0895-4356\(89\)90065-6](https://doi.org/10.1016/0895-4356(89)90065-6)
 43. Agrawal A, Galwankar S, Kapil V, Coronado V, Basavaraju S, McGuire L, et al. Epidemiology and clinical characteristics of traumatic brain injuries in a rural setting in Maharashtra, India. 2007–2009. *International Journal of Critical Illness and Injury Science*. 2012; 2(3): 167. <https://doi.org/10.4103/2229-5151.100915>
 44. Bhalerao G, Kulkarni V, Kapoor D. Comparison of Motor Relearning Program and Bobath Approach in acute stroke rehabilitation. *Journal of Orthopaedics and Rehabilitation*. 2011; 1(1): 79–88.
 45. Samanamalee S, Sigera PC, De Silva AP, Thilakasiri K, Rashan A, Wadanambi S, et al. Traumatic brain injury (TBI) outcomes in an LMIC tertiary care centre and performance of trauma scores. *BMC Anesthesiology*. 2018; 18(1): 4. <https://doi.org/10.1186/s12871-017-0463-7>
 46. Rowland TJ, Cooke DM, Gustafsson LA. Role of occupational therapy after stroke. *Annals of Indian Academy of Neurology*. 2008; 11(5): 99–107
 47. Skubik-Peplaski C, Custer M, Powell E, Westgate PM, Sawaki L. Comparing Occupation-Based and Repetitive Task Practice Interventions for Optimal Stroke Recovery: A Pilot Randomized Trial. *Physical & Occupational Therapy In Geriatrics*. 2017; 35(3–4): 156–168. <https://doi.org/10.1080/02703181.2017.1342734>
 48. Giuffrida CG, Demery JA, Reyes LR, Lebowitz BK, Hanlon RE. Functional Skill Learning in Men With Traumatic Brain Injury. *The American Journal Of Occupational Therapy*. 2009; 63(4): 398–407. <https://doi.org/10.5014/ajot.63.4.398>
 49. Shepherd RB. Exercise and training to optimize functional motor performance in stroke: Driving neural reorganization? *Neural Plasticity*. 2001; 8(1–2): 121–129. <https://doi.org/10.1155/np.2001.121>
 50. Huseynsinoglu BE, Ozdinciler AR, Krespi Y. Bobath Concept versus constraint-induced movement therapy to improve arm functional recovery in stroke patients: A randomized controlled trial. *Clinical Rehabilitation*. 2012; 26(8): 705–715. <https://doi.org/10.1177/0269215511431903>
 51. Hiragami S, Inoue Y, Harada K. Minimal clinically important difference for the Fugl-Meyer assessment of the upper extremity in convalescent stroke patients with moderate to severe hemiparesis. *Journal of Physical Therapy Science*. 2019; 31(11): 917–921: <https://doi.org/10.1589/jpts.31.917>
 52. Platz T, Winter T, Müller N, Pinkowski C, Eickhof C, Mauritz KH. Arm ability training for stroke and traumatic brain injury patients

- with mild arm paresis: A single-blind, randomized, controlled trial. *Archives of Physical Medicine and Rehabilitation*. 2001; 82(7): 961–968. <https://doi.org/10.1053/apmr.2001.23982>
53. Khan F, Baguley IJ, Cameron ID. Rehabilitation after traumatic brain injury. *Medical Journal of Australia*. 2003; 178(6): 290–295.
54. Lamontagne ME, Gagnon C, Allaire AS, Noreau L. Effect of rehabilitation length of stay on outcomes in individuals with traumatic brain injury or spinal cord injury: a systematic review protocol. *Systematic reviews*. 2013; 2: Article 59. <https://doi.org/10.1186/2046-4053-2-59>
55. Walsh K. Management of shoulder pain in patients with stroke. *Postgraduate Medical Journal*. 2001; 77(912): 645–649. <https://doi.org/10.1136/pmj.77.912.645>
56. Zhu XL, Poon WS, Chan CCH, Chan SSH. Does intensive rehabilitation improve the functional outcome of patients with traumatic brain injury (TBI)? A randomized controlled trial. *Brain Injury*. 2007; 21(7): 681–690. <https://doi.org/10.1080/02699050701468941>
57. Seneviratne C, Reimer M. Neurodevelopmental treatment stroke rehabilitation: a critique and extension for neuroscience nursing practice. *Axone*. 2004; 26(2): 13–20.

ROLES OF AUTHORS

Jackson Nowa is a post graduate student at the Wits University who was responsible for conceptualization of the research including connecting ideas and formulation of research goals. He was responsible for data curation including management of raw data and application of statistical analysis. He managed the investigation process including seeking different approvals from the university as well as hospitals and the ministry of Health and Social Services in Namibia, and the initial writing of the original draft and corrected versions.

Denise Franzsen was a Lead supervisor, and was responsible for supervision of the research including research planning, execution and mentorship of the post graduate student. She performed validation which involved verification of research components. She was responsible for reviewing and editing write ups, critical review, commentary and revision pre and post publication stages.

Dineo Thupae was the co-supervisor, and was responsible for supervision of the research including research planning, execution and mentorship of the post graduate student. She performed validation which involved verification of research components. She was responsible for reviewing and editing write ups, critical review, commentary and revision pre and post publication stages.

Corresponding Author

*Jackson Nowa

Email: jsn547644@gmail.com